



[Go to Product page](#)

Datasheet for ABIN1700973
anti-PBX4 antibody (AA 201-320) (Biotin)

Overview

Quantity:	100 µL
Target:	PBX4
Binding Specificity:	AA 201-320
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PBX4 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PBX4
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PBX4
Alternative Name:	PBX4 (PBX4 Products)
Background:	Synonyms: 2410015M21Rik, AI429113, AW045266, Edg4, Endothelial differentiation,

Target Details

lysophosphatidic acid G protein coupled receptor 4, Homeobox protein PBX4, OTTMUSP00000045221, PBX4, PBX4_HUMAN, pre-B-cell leukemia homeobox 4, Pre-B-cell leukemia transcription factor 4.

Background: Pbx 1, 2, 3 and 4 are members of the TALE (three amino acid loop extension) family of homeodomain-containing proteins. Human pre-B cell acute leukemias are frequently associated with a t(1,19)(q23,p13.3) chromosomal rearrangement, which creates a chimeric gene encoding a fusion between the E2A and Pbx 1 gene products. Pbx 2 and Pbx 3 share 92 % and 94 % respective identities with Pbx 1 over a 266 amino acid region flanking their homeobox domains, while all three proteins are quite divergent at their amino- and carboxy-termini. Two forms of Pbx 1 and Pbx 3 each differ primarily in their carboxy-termini and result from alternative mRNA splicing. Unlike other homeotic selector genes which are expressed transiently during development and differentiation, Pbx gene transcripts are ubiquitously expressed in both fetal and adult tissues and cell lines. Additionally, Pbx 2 and Pbx 3 transcripts are detected in lymphoid cells, which do not express Pbx 1. Pbx 4 expression is confined to the testis, especially to spermatocytes in the pachytene stage of the first meiotic prophase.

Gene ID: 80714

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months