



[Go to Product page](#)

Datasheet for ABIN1871737  
**anti-TNKS2 antibody (AA 959-1166)**

2 Images

Overview

Quantity:	100 µL
Target:	TNKS2
Binding Specificity:	AA 959-1166
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNKS2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Tankyrase 2 (TNKS2)
Immunogen:	Recombinant Tankyrase 2 (TNKS2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TNKS2. It has been selected for its ability to recognize TNKS2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details

---

Target:	TNKS2
Alternative Name:	Tankyrase 2 ( <a href="#">TNKS2 Products</a> )
Background:	PARP5B, PARP5C, TANK2, TNKL, Tankyrase-like protein, ADP-ribosyltransferase diphtheria toxin-like 6, TRF1-Interacting Ankyrin-Related ADP-Ribose Polymerase 2

## Application Details


---

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Concentration:	Lot specific
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



### Immunohistochemistry

**Image 1.** Used in DAB staining on formalin fixed paraffin-embedded liver tissue



### Western Blotting

**Image 2.** Figure. Western Blot; Sample: Recombinant TNKS2, Human.