



[Go to Product page](#)

Datasheet for ABIN1881847

anti-STAT1 antibody (C-Term)

6 Images

4 Publications

Overview

Quantity:	400 µL
Target:	STAT1
Binding Specificity:	AA 717-745, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF), Flow Cytometry (FACS)

Product Details

Immunogen:	This STAT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 717-745 amino acids from the C-terminal region of human STAT1.
Clone:	RB41406
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	STAT1
Alternative Name:	STAT1 (STAT1 Products)

Target Details

Background:	The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq].
Molecular Weight:	87335
NCBI Accession:	NP_009330 , NP_644671
UniProt:	P42224
Pathways:	JAK-STAT Signaling , RTK Signaling , Interferon-gamma Pathway , Response to Growth Hormone Stimulus , Cellular Response to Molecule of Bacterial Origin , Positive Regulation of Endopeptidase Activity , Hepatitis C , CXCR4-mediated Signaling Events

Application Details

Application Notes:	IF: 1:25. WB: 1:1000. IHC-P-Leica: 1:1000. IHC-P-Leica: 1:500. IHC-P-Leica: 1:500. FC: 1:25
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Expiry Date:	6 months

Publications

Product cited in:	Dai, Liu, Liu, Zhang, Wang, Jin, Qian, Wang, Zhao, Wu, Xiong, Chang, Sun, Yang, Hoffman, Liu: "Anti-metastatic Efficacy of Traditional Chinese Medicine (TCM) Ginsenoside Conjugated to a
-------------------	---


VEGFR-3 Antibody on Human Gastric Cancer in an Orthotopic Mouse Model." in: **Anticancer research**, Vol. 37, Issue 3, pp. 979-986, (2017) ([PubMed](#)).

Irrthum, Karkkainen, Devriendt, Alitalo, Vikkula: "Congenital hereditary lymphedema caused by a mutation that inactivates VEGFR3 tyrosine kinase." in: **American journal of human genetics**, Vol. 67, Issue 2, pp. 295-301, (2000) ([PubMed](#)).

Galland, Karamysheva, Pebusque, Borg, Rottapel, Dubreuil, Rosnet, Birnbaum: "The FLT4 gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor." in: **Oncogene**, Vol. 8, Issue 5, pp. 1233-40, (1993) ([PubMed](#)).


Pajusola, Aprelikova, Korhonen, Kaipainen, Pertovaara, Alitalo, Alitalo: "FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines." in: **Cancer research**, Vol. 52, Issue 20, pp. 5738-43, (1992) ([PubMed](#)).

Galland, Karamysheva, Mattei, Rosnet, Marchetto, Birnbaum: "Chromosomal localization of FLT4, a novel receptor-type tyrosine kinase gene." in: **Genomics**, Vol. 13, Issue 2, pp. 475-8, (1992) ([PubMed](#)).




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of paraffin-embedded human brain tissue using (ABIN1881847 and ABIN2838812) performed on the Leica® BOND RXm. Samples were incubated with primary Antibody (1/500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunofluorescence

Image 2. Immunofluorescent analysis of 4 % paraformaldehyde-fixed, 0.1 % Triton X-100 permeabilized HeLa cells labeling STAT1 with (ABIN1881847 and ABIN2838812) at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-Rabbit IgG secondary antibody at 1/200 dilution (green). Immunofluorescence image showing Nucleus and Weak Cytoplasm staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin(red). The nuclear counter stain is DI (blue).



Western Blotting

Image 3. All lanes : Anti-STAT1 Antibody (C-term) at 1:1000 dilution Lane 1: Jurkat whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: Daudi whole cell lysate Lane 5: HeLa whole cell lysate Lane 5: MCF-7 whole cell lysate Lane 5: HT-29 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 87 kDa Blocking/Dilution buffer: 5 % NFDN/TBST.

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN1881847.