



[Go to Product page](#)

Datasheet for ABIN1991794  
**anti-CETN2 antibody (AA 1-172) (FITC)**

### Overview

Quantity:	100 µg
Target:	CETN2
Binding Specificity:	AA 1-172
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CETN2 antibody is conjugated to FITC
Application:	Western Blotting (WB), ELISA

### Product Details

Immunogen:	Recombinant Human Centrin-2 protein (1-172AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

### Target Details

Target:	CETN2
Alternative Name:	CETN2 ( <a href="#">CETN2 Products</a> )
Background:	Background: Plays a fundamental role in microtubule organizing center structure and function. Required for centriole duplication and correct spindle formation. Has a role in regulating

## Target Details

---

cytokinesis and genome stability via cooperation with CALM1 and CCP110. Involved in global genome nucleotide excision repair (GG-NER) by acting as component of the XPC complex. Cooperatively with RAD23B appears to stabilize XPC. In vitro, stimulates DNA binding of the XPC:RAD23B dimer. The XPC complex is proposed to represent the first factor bound at the sites of DNA damage and together with other core recognition factors, XPA, RPA and the TFIIH complex, is part of the pre-incision (or initial recognition) complex. The XPC complex recognizes a wide spectrum of damaged DNA characterized by distortions of the DNA helix such as single-stranded loops, mismatched bubbles or single-stranded overhangs. The orientation of XPC complex binding appears to be crucial for inducing a productive NER. XPC complex is proposed to recognize and to interact with unpaired bases on the undamaged DNA strand which is followed by recruitment of the TFIIH complex and subsequent scanning for lesions in the opposite strand in a 5'-to-3' direction by the NER machinery. Cyclobutane pyrimidine dimers (CPDs) which are formed upon UV-induced DNA damage escape detection by the XPC complex due to a low degree of structural perturbation. Instead they are detected by the UV-DDB complex which in turn recruits and cooperates with the XPC complex in the respective DNA repair.

Aliases: 20kD calcium binding protein antibody, CALT antibody, caltractin antibody, Caltractin isoform 1 antibody, CEN2 antibody, centrin antibody, centrin, EF hand protein, 2 antibody, Centrin-2 antibody, Centrin2 antibody, CETN2 antibody, CETN2\_HUMAN antibody, EF hand protein 2 antibody, EF-hand protein antibody

---

UniProt: [P41208](#)

---

Pathways: [DNA Damage Repair, M Phase](#)

## Application Details

---

Application Notes: Optimal working concentration should be determined by the investigator.

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin™: a POISONOUS AND HAZARDOUS SUBSTANCE, which should

---

## Handling

---

be handled by trained staff only.

---

Handling Advice: Avoid repeated freeze.

---

Storage: -20 °C,-80 °C

---

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.