



Datasheet for ABIN2704706 anti-KRT20 antibody (C-Term)

[Go to Product page](#)

3 Images

Overview

Quantity: 200 µL

Target: KRT20

Binding Specificity: C-Term

Reactivity: Human, Mouse, Rat, Pig, Cow

Host: Rabbit

Clonality: Polyclonal

Conjugate: This KRT20 antibody is un-conjugated

Application: Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),
Immunochromatography (IC)

Product Details

Immunogen: KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 20.

Specificity: Recognizes endogenous levels of Cytokeratin 20 protein.

Characteristics: Rabbit polyclonal antibody to Cytokeratin 20

Purification: The antibody was purified by immunogen affinity chromatography.

Target Details

Target: KRT20

Alternative Name: Cytokeratin 20 ([KRT20 Products](#))

Target Details

Background:	Keratin, type I cytoskeletal 20, Cytokeratin-20, CK-20, Keratin-20, K20, Protein IT
Gene ID:	54474, 66809, 286912
UniProt:	P35900 , Q9D312 , P25030


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence


Image 1. Immunofluorescent analysis of Cytokeratin 20 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI



was used to stain the cell nuclei (blue).

Immunohistochemistry

Image 2. Immunohistochemical analysis of Cytokeratin 20 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of Cytokeratin 20 expression in A549 (A), NS-1 (B), mouse kidney (C), rat kidney (D) whole cell lysates.