



Datasheet for ABIN2706306

anti-HDAC5 antibody (pSer498)

3 Images



[Go to Product page](#)

Overview

Quantity: 200 µL

Target: HDAC5

Binding Specificity: pSer498

Reactivity: Human, Mouse, Rat, Cow, Pig

Host: Rabbit

Clonality: Polyclonal

Conjugate: This HDAC5 antibody is un-conjugated

Application: Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),
Immunochromatography (IC)

Product Details

Immunogen: KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone Deacetylase 5.

Specificity: Recognizes endogenous levels of Histone Deacetylase 5 (pS498) protein.

Characteristics: Rabbit polyclonal antibody to Histone Deacetylase 5 (pS498)

Purification: The antibody was purified by immunogen affinity chromatography.

Target Details

Target: HDAC5

Alternative Name: Histone Deacetylase 5 ([HDAC5 Products](#))

Target Details


Background:	KIAA0600, Histone deacetylase 5, HD5, Antigen NY-CO-9
Gene ID:	10014
UniProt:	Q9UQL6 , Q9Z2V6
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development, Monocarboxylic Acid Catabolic Process

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




Immunofluorescence

Image 1. Immunofluorescent analysis of Histone Deacetylase 5 (pS498) staining in Raw264.7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemistry

Image 2. Immunohistochemical analysis of Histone Deacetylase 5 (pS498) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of Histone Deacetylase 5 (pS498) expression in HEK293T LPS-treated (A), Raw264.7 TNFa-treated (B), rat kidney (C) whole cell lysates.