



Datasheet for ABIN3093042

## IFI16 Protein (AA 1-785) (Strep Tag)



[Go to Product page](#)

### 1 Image

#### Overview

Quantity:	1 mg
Target:	IFI16
Protein Characteristics:	AA 1-785
Origin:	Human
Source:	Tobacco ( <i>Nicotiana tabacum</i> )
Protein Type:	Recombinant
Purification tag / Conjugate:	This IFI16 protein is labelled with Strep Tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA

#### Product Details

Sequence: MGKKYKNIVL LKGLEVINDY HFRMVKSLLS NDLKLNLMR EEYDKIQIAD LMEEKFRGDA  
GLGKLIKIFE DIPTLEDLAE TLKKEKLVK GPALSRKRKK EVDATSPAPS TSSTVKTEGA  
EATPGAQKRK KSTKEKAGPK GSKVSEEQTQ PPSAGAGMS TAMGRSPSPK TSLSAPPNSS  
STENPKTVAK CQVTPRRNVL QKRPVIVKVL STTKPFYET PEMEKKIMFH ATVATQTQFF  
HVKVLNTSLK EKFNKGI III ISDYLEYDSL LEVNEESTVS EAGPNQTFEV PNKIINRAKE  
TLKIDILHKQ ASGNIVGVF MLHKKTVNQK TTIYEIQDDR GKMDVVG TGQ CHNIPCEEGD  
KLQLFCFRLR KKNQMSKLIS EMHSFIQIKK KTNPRNNDPK SMKLPQEQRQ LPYPSEASTT  
FPESHLRTPQ MPPTTPSSSF FTKKSED TIS KMNDFMRMQ I LKEGSHFPGP FM TSIGPAES  
HPHTPQMPPS TPSSSFLTTK SEDTISKMND FMRMQILKEG SHFPGPFMTS IGPAESHPH T  
PQMPPSTPSS SFLTTLK PRL KTEPEEVSIE DSAQSDLKEV MVLNATESFV YEPKEQKKMF  
HATVATENEV FRVKVFNIDL KEKFTP KII AIANYVCRNG FLEVYPFTLV ADVNADR NME  
IPKGLIR SAS VTPKINQLCS QTKGSFVNGV FEVHKKNVRG EFTYIEIQDN TGKMEVVVHG

RLTTINCEEG DKLKLTQFEL APKSGNTGEL RSVIHSIKV IKTRKNKKDI LNPDSSMETS PDFFF

**Sequence without tag. The proposed Strep-Tag is based on experience s with the expression system, a different complexity of the protein could make another tag necessary. In case you have a special request, please contact us.**

---

### Characteristics:

#### Key Benefits:

- Made in Germany - from design to production - by highly experienced protein experts.
- Protein expressed with ALiCE® and purified by multi-step, protein-specific process to ensure correct folding and modification.
- These proteins are normally active (enzymatically functional) as our customers have reported (not tested by us and not guaranteed).
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a **made-to-order protein** and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our **made-to-order proteins** in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

#### Expression System:

- ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.
- During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

#### Concentration:

- The concentration of our recombinant proteins is measured using the absorbance at 280nm.
- The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.
- We use the Expasy's ProtParam tool to determine the absorption coefficient of each protein.

### Purification:

Two step purification of proteins expressed in Almost Living Cell-Free Expression System

---

## Product Details

---

(ALiCE®):

1. In a first purification step, the protein is purified from the cleared cell lysate using StrepTag capture material. Eluate fractions are analyzed by SDS-PAGE.
2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.

---

Purity: >80 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

---

Endotoxin Level: Low Endotoxin less than 1 EU/mg (< 0.1 ng/mg)

---

Grade: Crystallography grade

## Target Details

---

Target: IFI16

---

Alternative Name: IFI16 ([IFI16 Products](#))

---

Background: Gamma-interferon-inducible protein 16 (I $\gamma$ I-16) (Interferon-inducible myeloid differentiation transcriptional activator),FUNCTION: Binds double-stranded DNA. Binds preferentially to supercoiled DNA and cruciform DNA structures. Seems to be involved in transcriptional regulation. May function as a transcriptional repressor. Could have a role in the regulation of hematopoietic differentiation through activation of unknown target genes. Controls cellular proliferation by modulating the functions of cell cycle regulatory factors including p53/TP53 and the retinoblastoma protein. May be involved in TP53-mediated transcriptional activation by enhancing TP53 sequence-specific DNA binding and modulating TP53 phosphorylation status. Seems to be involved in energy-level-dependent activation of the ATM/ AMPK/TP53 pathway coupled to regulation of autophagy. May be involved in regulation of TP53-mediated cell death also involving BRCA1. May be involved in the senescence of prostate epithelial cells. Involved in innate immune response by recognizing viral dsDNA in the cytosol and probably in the nucleus. After binding to viral DNA in the cytoplasm recruits TMEM173/STING and mediates the induction of IFN-beta. Has anti-inflammatory activity and inhibits the activation of the AIM2 inflammasome, probably via association with AIM2. Proposed to bind viral DNA in the nucleus, such as of Kaposi's sarcoma-associated herpesvirus, and to induce the formation of nuclear caspase-1-activating inflammasome formation via association with PYCARD. Inhibits replication of herpesviruses such as human cytomegalovirus (HCMV) probably by interfering with promoter recruitment of members of the Sp1 family of transcription factors. Necessary to activate the IRF3 signaling cascade during human herpes simplex virus 1 (HHV-1) infection and promotes the assembly of heterochromatin on herpesviral DNA and inhibition of viral

## Target Details

---

immediate-early gene expression and replication. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer. {ECO:0000269|PubMed:11146555, ECO:0000269|PubMed:12894224, ECO:0000269|PubMed:14654789, ECO:0000269|PubMed:20890285, ECO:0000269|PubMed:21573174, ECO:0000269|PubMed:21575908, ECO:0000269|PubMed:22046441, ECO:0000269|PubMed:22291595, ECO:0000269|PubMed:23027953, ECO:0000269|PubMed:24198334, ECO:0000269|PubMed:24413532, ECO:0000269|PubMed:9642285}., FUNCTION: [Isoform IFI16-beta]: Isoform that specifically inhibits the AIM2 inflammasome (PubMed:30104205). Binds double-stranded DNA (dsDNA) in the cytoplasm, impeding its detection by AIM2 (PubMed:30104205). Also prevents the interaction between AIM2 and PYCARD/ASC via its interaction with AIM2, thereby inhibiting assembly of the AIM2 inflammasome (PubMed:30104205). This isoform also weakly induce production of type I interferon-beta (IFNB1) via its interaction with STING1 (PubMed:30104205). {ECO:0000269|PubMed:30104205}.

---

Molecular Weight: 88.3 kDa

---

UniProt: [Q16666](#)

---

Pathways: [Activation of Innate immune Response](#), [Positive Regulation of Endopeptidase Activity](#), [Autophagy](#), [Inflammasome](#)

## Application Details

---

Application Notes: In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.

Comment: ALiCE®, our Almost Living Cell-Free Expression System is based on a lysate obtained from *Nicotiana tabacum* c.v.. This contains all the protein expression machinery needed to produce even the most difficult-to-express proteins, including those that require post-translational modifications.

During lysate production, the cell wall and other cellular components that are not required for protein production are removed, leaving only the protein production machinery and the mitochondria to drive the reaction. During our lysate completion steps, the additional components needed for protein production (amino acids, cofactors, etc.) are added to produce something that functions like a cell, but without the constraints of a living system - all that's needed is the DNA that codes for the desired protein!

---

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Buffer:	The buffer composition is at the discretion of the manufacturer. If you have a special request, please contact us.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.
Expiry Date:	Unlimited (if stored properly)

## Images

---



**Image 1.** „Crystallography Grade“ protein due to multi-step, protein-specific purification process