



Datasheet for ABIN3134784

Serotonin Receptor 2B Protein (HTR2B) (AA 1-479) (rho-1D4 tag)



[Go to Product page](#)

1 Image

Overview

Quantity:	1 mg
Target:	Serotonin Receptor 2B (HTR2B)
Protein Characteristics:	AA 1-479
Origin:	Mouse
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Serotonin Receptor 2B protein is labelled with rho-1D4 tag.
Application:	ELISA, Western Blotting (WB), Crystallization (Crys), SDS-PAGE (SDS)

Product Details

Sequence: MASSYKMSEQ STTSEHILQK TCDHLILTNR SGLETDSVAE EMKQTVEGQG HTVHWAALLI
LAVIPIPTIGG NILVILAVAL EKRLQYATNY FLMSLAIADL LVGLFVMPIA LLTIMFEAIW
PLPLALCPAW LFLDVLFSTA SIMHLCAISL DRYIAIKKPI QANQCNSRAT AFIKITVVWL ISIGIAIPVP
IKGIETDVIN PHNVTCELT KDRFGSFMVFG SLAAFFAPLT IMVVTYFLTI HTLQKKAYLV
KNKPPQRLTR WTVPTVFLRE DSSFSSPEKV AMLDGSHRD K ILPNSSDETL MRRMSSVGKR
SAQTISNEQR ASKALGVVFF LFLLMWCPFF ITNLTLACD SCNQTTLKTL LEIFVWIGYV
SSGVNPLIYT LFNKTFR EAF GRYITCNYRA TKSVKALRK F SSTLCFGNSM VENSKFFT KH
GIRNGINPAM YQSPMRLRSS TIQSSSIILL DTLLTENDGD KAEEQVSYI

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

Characteristics:

- Made in Germany - from design to production - by highly experienced protein experts.
- Mouse Htr2b Protein (raised in Insect Cells) purified by multi-step, protein-specific process to

Product Details

- ensure crystallization grade.
- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made-to-order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom-made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification: Three step purification of membrane proteins expressed in baculovirus infected SF9 insect cells:

1. Membrane proteins are fractionated by ultracentrifugation and subsequently solubilized with different detergents (detergent screen). Samples are analyzed by Western blot.
2. The best performing detergent is used for solubilization and the proteins are purified via their rho1D4 tag via two rho1D4 antibody columns: one DTT resistant, the other one not. Eluate fractions are analyzed by Western blot.
3. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatograph. Eluate fractions are analyzed by SDS-PAGE and Western blot.

Purity: >95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.

Sterility: 0.22 µm filtered

Endotoxin Level: Protein is endotoxin-free.

Grade: Crystallography grade

Target Details

Target:	Serotonin Receptor 2B (HTR2B)
Alternative Name:	Htr2b (HTR2B Products)
Background:	<p>G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various ergot alkaloid derivatives and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways.</p> <p>Signaling activates a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and down-stream signaling cascades and promotes the release of Ca(2+) ions from intracellular stores. Plays a role in the regulation of dopamine and 5-hydroxytryptamine release, 5-hydroxytryptamine uptake and in the regulation of extracellular dopamine and 5-hydroxytryptamine levels, and thereby affects neural activity.</p> <p>May play a role in the perception of pain. Plays a role in the regulation of behavior, including impulsive behavior. Required for normal proliferation of embryonic cardiac myocytes and normal heart development. Protects cardiomyocytes against apoptosis. Plays a role in the adaptation of pulmonary arteries to chronic hypoxia. Plays a role in vasoconstriction. Required for normal osteoblast function and proliferation, and for maintaining normal bone density.</p> <p>Required for normal proliferation of the interstitial cells of Cajal in the intestine.</p> <p>{ECO:0000269 PubMed:10944220, ECO:0000269 PubMed:11413089, ECO:0000269 PubMed:12244304, ECO:0000269 PubMed:12738797, ECO:0000269 PubMed:1426253, ECO:0000269 PubMed:16940156, ECO:0000269 PubMed:17846081, ECO:0000269 PubMed:18337424, ECO:0000269 PubMed:19941613, ECO:0000269 PubMed:23346101}.</p>
Molecular Weight:	54.8 kDa Including tag.
UniProt:	Q02152
Pathways:	JAK-STAT Signaling , Inositol Metabolic Process , Regulation of G-Protein Coupled Receptor Protein Signaling , Regulation of Carbohydrate Metabolic Process

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	Protein has not been tested for activity yet. In cases in which it is highly likely that the

Application Details

recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: 100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: -80 °C

Storage Comment: Store at -80°C.

Expiry Date: Unlimited (if stored properly)

Images



Image 1. „Crystallography Grade“ protein due to multi-step, protein-specific purification process