



Datasheet for ABIN6258844

anti-HNRPLL antibody (Internal Region)

3 Images



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HNRPLL
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRPLL antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human HNRPLL, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	HNRPLL Antibody detects endogenous levels of total HNRPLL.
Predicted Reactivity:	Pig, Bovine, Horse, Sheep, Rabbit, Dog, Xenopus

Target Details

Target:	HNRPLL
---------	--------

Target Details

Alternative Name:	HNRNPLL (HNRPLL Products)
Background:	Description: RNA-binding protein that functions as regulator of alternative splicing for multiple target mRNAs, including PTPRC/CD45 and STAT5A. Required for alternative splicing of PTPRC. Gene: HNRNPLL
Molecular Weight:	60 kDa
Gene ID:	92906
UniProt:	Q8WVV9

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Immunofluorescence (fixed cells)

Image 1. ABIN6275184 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 2. Western blot analysis of extracts from mouse brain, using HNRPLL Antibody.



Western Blotting

Image 3. Western blot analysis of extracts from COLO cells using HNRPLL antibody.