



[Go to Product page](#)

Datasheet for ABIN6262315

## anti-HNRNPA1 antibody (Internal Region)

### 4 Images

#### Overview

Quantity:	100 µL
Target:	HNRNPA1
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPA1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

#### Product Details

Immunogen:	A synthesized peptide derived from human hnRNP A1, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	HnRNP A1 Antibody detects endogenous levels of total hnRNP A1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

#### Target Details

Target:	HNRNPA1
---------	---------

## Target Details

---

Alternative Name:	HNRNPA1 ( <a href="#">HNRNPA1 Products</a> )
Background:	Description: Involved in the packaging of pre-mRNA into hnRNP particles, transport of poly(A) mRNA from the nucleus to the cytoplasm and may modulate splice site selection (PubMed:17371836). May bind to specific miRNA hairpins (PubMed:28431233). Gene: HNRNPA1
Molecular Weight:	39 kDa
Gene ID:	3178
UniProt:	<a href="#">P09651</a>

## Application Details


---

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




### Immunohistochemistry

**Image 1.** ABIN6268816 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



### Western Blotting

**Image 2.** Western blot analysis of hnRNP A1 expression in 293 cell extract



### Immunofluorescence (fixed cells)

**Image 3.** ABIN6268816 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN6262315.