



[Go to Product page](#)

Datasheet for ABIN6265092

anti-SIRT3 antibody (C-Term)

3 Images

1 Publication

Overview

Quantity:	100 µL
Target:	SIRT3
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SIRT3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human SIRT3, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	SIRT3 Antibody detects endogenous levels of total SIRT3.
Predicted Reactivity:	Sheep,Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	SIRT3
---------	-------

Target Details

Alternative Name: SIRT3 ([SIRT3 Products](#))

Background: Description: NAD-dependent protein deacetylase (PubMed:12186850, PubMed:12374852, PubMed:16788062, PubMed:18680753, PubMed:18794531, PubMed:23283301, PubMed:24121500, PubMed:24252090, PubMed:19535340). Activates or deactivates mitochondrial target proteins by deacetylating key lysine residues (PubMed:12186850, PubMed:12374852, PubMed:16788062, PubMed:18680753, PubMed:18794531, PubMed:23283301, PubMed:24121500, PubMed:24252090). Known targets include ACSS1, IDH, GDH, SOD2, PDHA1, LCAD, SDHA and the ATP synthase subunit ATP5O (PubMed:16788062, PubMed:18680753, PubMed:24121500, PubMed:24252090, PubMed:19535340). Contributes to the regulation of the cellular energy metabolism (PubMed:24252090). Important for regulating tissue-specific ATP levels (PubMed:18794531). In response to metabolic stress, deacetylates transcription factor FOXO3 and recruits FOXO3 and mitochondrial RNA polymerase POLRMT to mtDNA to promote mtDNA transcription (PubMed:23283301).

Gene: SIRT3

Molecular Weight: 29 kDa

Gene ID: 23410

UniProt: [Q9NTG7](#)

Application Details

Application Notes: WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20 °C. Stable for 12 months from date of receipt.

Expiry Date: 12 months

Publications


Product cited in: Wu, Huang, Jiao, Ding, Zhang, Chen, Wang, Li, Huo: "Olaquinox disrupts tight junction integrity and cytoskeleton architecture in mouse Sertoli cells." in: **Oncotarget**, Vol. 8, Issue 51, pp. 88630-88644, (2017) ([PubMed](#)).

Images




Immunofluorescence (fixed cells)

Image 1. ABIN6268681 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 2. Western blot analysis of SIRT3 expression in Mouse Heart lysate, The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 3. ABIN6268681 at 1/100 staining Mouse testis tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22;ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary