



[Go to Product page](#)

Datasheet for ABIN6266376
anti-PGC antibody (Internal Region)

3 Images

Overview

Quantity:	100 µL
Target:	PGC
Binding Specificity:	Internal Region
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PGC antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human PGC, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	PGC Antibody detects endogenous levels of total PGC.
Predicted Reactivity:	Bovine, Sheep, Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	PGC
---------	-----

Target Details


Alternative Name:	PGC (PGC Products)
Background:	Description: Hydrolyzes a variety of proteins. Gene: PGC
Molecular Weight:	42kDa
Gene ID:	5225
UniProt:	P20142

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:100, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts from hybridoma cells, using PGC Antibody. Lane 1 was treated with the blocking peptide.



Western Blotting

Image 2. Western blot analysis of extracts of 823, using PGC antibody. The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 3. ABIN6277340 at 1/100 staining Human uterus tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary