



[Go to Product page](#)

Datasheet for ABIN657511
anti-RHBDF2 antibody (N-Term)

5 Images

Overview

| | |
|----------------------|--|
| Quantity: | 200 µL |
| Target: | RHBDF2 |
| Binding Specificity: | AA 80-109, N-Term |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| | |
|---------------|---|
| Immunogen: | This RHBDF2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 80-109 amino acids from the N-terminal region of human RHBDF2. |
| Clone: | RB32834 |
| Isotype: | IgG |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|---|
| Target: | RHBDF2 |
| Alternative Name: | RHBDF2 (RHBDF2 Products) |
| Background: | Not expected to have protease activity (By similarity). |
| Molecular Weight: | 96686 |

Target Details

| | |
|-----------------|--|
| Gene ID: | 79651 |
| NCBI Accession: | NP_001005498 , NP_078875 |
| UniProt: | Q6PJF5 |
| Pathways: | Growth Factor Binding |


Application Details

| | |
|--------------------|--|
| Application Notes: | WB: 1:2000. WB: 1:2000. WB: 1:1000. IHC-P: 1:25. IHC-P: 1:25 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | RHBDF2 Antibody (N-term) can be refrigerated at 2-8 °C for up to 6 months. For long term storage, place the at -20 °C. |
| Expiry Date: | 6 months |


Images



Immunohistochemistry (Paraffin-embedded Sections)


Image 1. (ABIN657511 and ABIN2846535) staining RHBDF2 in human duodenum tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3 % BSA for 0.5 hour at room temperature, antigen retrieval was by heat mediation with a citrate buffer (pH 6). Samples were incubated with primary antibody (1/25) for 1 hours at 37 °C. A undiluted

biotinylated goat polyvalent antibody was used as the secondary antibody.



Western Blotting

Image 2. Anti-RHBDF2 Antibody (N-term) at 1:2000 dilution + Mouse lung tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 97 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.



Western Blotting

Image 3. Anti-RHBDF2 Antibody (N-term) at 1:2000 dilution + HACAT whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 97 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN657511.