



## Datasheet for ABIN749888 anti-TREML2 antibody (AA 65-150)



[Go to Product page](#)

### Overview

Quantity:	100 µL
Target:	TREML2
Binding Specificity:	AA 65-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TREML2 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TREML2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

### Target Details

Target:	TREML2
Alternative Name:	<a href="#">TREML2 (TREML2 Products)</a>

## Target Details

---

Background:	Synonyms: C6orf76, Chromosome 6 open reading frame 76, FLJ13693, MGC149715, MGC149716, TLT 2, TLT2, TREM like transcript 2, Trem like transcript 2 protein, Triggering receptor expressed on myeloid cells like 2, Triggering receptor expressed on myeloid cells like protein 2, TRML2_HUMAN.  Background: TREML2 is expressed throughout B cell development in addition to being expressed on macrophages and neutrophils and is the only TREM molecule to be found on lymphocytes. TREML2 is expressed on B lineage cells early in development, and the highest level of expression is detected on those mature peripheral B cell subpopulations that are involved in the initial humoral immune response against bacterial pathogens. TREML2 is unique in that it lacks either the conserved transmembrane lysine residue or ITAM/ITIMs within its own cytoplasmic domain. Thus, TREML2 does not exhibit any of the features associated with classical tyrosine-based signaling. Monocytes in the bone marrow or peripheral blood do not express detectable levels of TREML2, but its expression is up-regulated in conjunction with differentiation into macrophages. TREML2 is present on neutrophils in the bone marrow as well as the periphery, and inflammatory stimuli result in a dramatic increase in the expression of TREML2 on these cells <i>in vivo</i> . TREML2 is a single-pass type I membrane protein, and it contains 1 Ig-like V-type (immunoglobulin-like) domain. It is a cell surface receptor that may play a role in the innate and adaptive immune response. TREML2 is located in a gene cluster on chromosome 6 with the single Ig variable (IgV) domain activating receptors TREM1 and TREM2, but it has distinct structural and functional properties.
-------------	---

---

Gene ID: 79865

UniProt: [Q5T2D2](#)

## Application Details

---

Application Notes: ELISA 1:500-1000  
IHC-P 1:200-400  
IHC-F 1:100-500  
IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

## Handling

---

Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months