



[Go to Product page](#)

Datasheet for ABIN886412

anti-NOD2 antibody (AA 251-350) (Alexa Fluor 488)

1 Image

Overview

Quantity:	100 µL
Target:	NOD2
Binding Specificity:	AA 251-350
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NOD2 antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CARD15
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NOD2
Alternative Name:	NOD2 (NOD2 Products)
Background:	Synonyms: ACUG, Arthrocutaneouveal granulomatosis, BLAU, CARD15, Caspase recruitment

Target Details

domain protein 15, Caspase recruitment domain-containing protein 15, IBD1, Inflammatory bowel disease protein 1, LRR containing protein, LRR-containing protein, Nod2, NOD2 protein, NOD2_HUMAN, NOD2B, nucleotide binding oligomerization domain 2, nucleotide-binding oligomerization domain 2, Nucleotide-binding oligomerization domain-containing protein 2, PSORAS1.

Background: Defects in NOD2 are the cause of sarcoidosis early-onset (EOS) . EOS is a form of sarcoidosis manifesting in children younger than 4 years of age. Sarcoidosis is an idiopathic, systemic, inflammatory disease characterized by the formation of immune granulomas in involved organs. Granulomas predominantly invade the lungs and the lymphatic system, but also skin, liver, spleen, eyes and other organs may be involved. Early-onset sarcoidosis is quite rare and has a distinct triad of skin, joint and eye disorders, without apparent pulmonary involvement. Compared with an asymptomatic and sometimes naturally disappearing course of the disease in older children, early-onset sarcoidosis is progressive and in many cases causes severe complications, such as blindness, joint destruction and visceral involvement.

Gene ID: 64127

UniProt: [Q9HC29](#)

Pathways: [Activation of Innate immune Response](#), [Cellular Response to Molecule of Bacterial Origin](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [Toll-Like Receptors Cascades](#), [Inflammasome](#)

Application Details

Application Notes: FCM 1:20-100

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling


handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months

Images



Flow Cytometry

Image 1. Raji probed with CARD15 Polyclonal Antibody, ALEXA FLUOR® 488 Conjugated (bs-7084R-A488) at 1:100 for 30 minutes compared to control cells (blue) and isotype control (orange).