



[Go to Product page](#)

Datasheet for ABIN928389

anti-FLASH antibody (N-Term)

1 Image

Overview

Quantity:	100 µL
Target:	FLASH (CASP8AP2)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FLASH antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	CASP8 AP2 antibody was raised in rabbit using the N terminal of CASP8 P2 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Dog (Canine)
Purification:	Purified

Target Details

Target:	FLASH (CASP8AP2)
Alternative Name:	CASP8AP2 (CASP8AP2 Products)
Background:	CASP8AP2 is highly similar to FLASH, a mouse apoptotic protein identified by its interaction with the death-effector domain (DED) of caspase 8. Studies of FLASH protein suggested that this protein may be a component of the death-inducing signaling complex that includes Fas receptor, Fas-binding adapter FADD, and caspase 8, and plays a regulatory role in Fas-mediated

Target Details

apoptosis. Synonyms: Polyclonal CASP8AP2 antibody, Anti-CASP8AP2 antibody, CASP8 associated protein 2 antibody, CED-4 antibody, FLASH antibody, FLJ11208 antibody, KIAA1315 antibody, RIP25 antibody.

Pathways: [Chromatin Binding](#), [Positive Regulation of Endopeptidase Activity](#)


Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	CASP8AP2 Blocking Peptide, catalog no. 33R-8602, is also available for use as a blocking control in assays to test for specificity of this CASP8AP2 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. CASP8AP2 antibody used at 0.2-1 ug/ml to detect target protein.