



Datasheet for ABIN950321
anti-AKR1C3 antibody (Middle Region)



[Go to Product page](#)

2 Images

Overview

Quantity:	0.4 mL
Target:	AKR1C3
Binding Specificity:	AA 114-143, Middle Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AKR1C3 antibody is un-conjugated
Application:	Western Blotting (WB), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	KLH conjugated synthetic peptide between 114-143 amino acids from the Central region of human AKR1C3
Isotype:	Ig Fraction
Specificity:	This antibody reacts to AKR1C3.
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Affinity chromatography on Protein A

Target Details

Target:	AKR1C3
Abstract:	AKR1C3 Products

Target Details

Background: This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reduction of prostaglandin (PG) D₂, PGH₂ and phenanthrenequinone (PQ), and the oxidation of 9 α ,11 β -PGF₂ to PGD₂. It may play an important role in the pathogenesis of allergic diseases such as asthma, and may also have a role in controlling cell growth and/or differentiation. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14. Synonyms: 17- β -hydroxysteroid dehydrogenase type 5, 2-dihydrobenzene-1, 2-diol dehydrogenase, 3- α -HSD type II, 3- α -HSD type II, 3- α -hydroxysteroid dehydrogenase type 2, Aldo-keto reductase family 1 member C3, Chlordecone reductase homolog HAKRb, Dihydrodiol dehydrogenase 3, Dihydrodiol dehydrogenase type I, HA1753, HSD17B5, KIAA0119, PGFS, Prostaglandin F synthase, Testosterone 17- β -dehydrogenase 5, Trans-1, brain

Gene ID: 8644

NCBI Accession: [NP_003730](#)

Pathways: [Retinoic Acid Receptor Signaling Pathway](#), [Steroid Hormone Biosynthesis](#), [Regulation of Hormone Metabolic Process](#), [Regulation of Hormone Biosynthetic Process](#), [C21-Steroid Hormone Metabolic Process](#), [Protein targeting to Nucleus](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.25 mg/mL

Buffer: PBS containing 0.09 % (W/V) sodium azide as preservative

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Handling Advice: Avoid repeated freezing and thawing.

Handling

Storage: 4 °C/-20 °C


Storage Comment: Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. AKR1C3 Antibody (Center) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of AKR1C3 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Western Blotting

Image 2. AKR1C3 Antibody (Center) western blot analysis in HeLa, HepG2 cell line lysates (35ug/lane). This demonstrates the AKR1C3 antibody detected the AKR1C3 protein (arrow).