



[Go to Product page](#)

Datasheet for ABIN950617

## anti-BCAT1 antibody (Middle Region)

### 2 Images

#### Overview

Quantity:	0.4 mL
Target:	BCAT1
Binding Specificity:	AA 88-115, Middle Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BCAT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

#### Product Details

Immunogen:	KLH conjugated synthetic peptide between 88-115 amino acids from the Central region of human BCAT1
Isotype:	Ig Fraction
Specificity:	This antibody reacts to BCAT1.
Cross-Reactivity (Details):	Species reactivity (tested):Human.
Purification:	Affinity chromatography on Protein A

#### Target Details

Target:	BCAT1
---------	-------

## Target Details

---

Alternative Name:	BCAT1 ( <a href="#">BCAT1 Products</a> )
Background:	<p>This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to branched-chain L-amino acids essential for cell growth. Two different clinical disorders have been attributed to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. As there is also a gene encoding a mitochondrial form of this enzyme, mutations in either gene may contribute to these disorders. Alternatively spliced transcript variants have been described. Synonyms: BCAT(c), BCT1, Branched-chain-amino-acid aminotransferase cytosolic, ECA39</p>
Gene ID:	586
NCBI Accession:	<a href="#">NP_001171562</a>

## Application Details


---

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	0.25 mg/mL
Buffer:	PBS containing 0.09 % (W/V) sodium azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** BCAT1 Antibody (Center) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the BCAT1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



### Western Blotting

**Image 2.** BCAT1 Antibody (Center) western blot analysis in Ramos cell line lysates (35µg/lane). This demonstrates the BCAT1 antibody detected the BCAT1 protein (arrow).